

Manipulating Relativistic Electrons with Intense Laser Pulses

Victor Malka^{1,2*}

¹*Laboratoire d'Optique Appliquée, CNRS, Ecole Polytechnique, ENSTA Paristech, Université Paris-Saclay, Palaiseau, France*

²*Weizmann Institute of Science, Rehovot, Israel*
**victor.malka@ensta.fr*

Laser Plasma Accelerators (LPA) rely on the control of the electronic motion with intense laser pulses [1]. The manipulation of electrons with intense laser pulses allows a fine mapping of the longitudinal and radial components of giant electric fields that can be therefore optimized for accelerating charged particle or for producing X rays. To illustrate the beauty of laser plasma accelerators I will show, how by changing the density profile of the gas target, one can improve the quality of the electron beam, its stability [2] and its energy gain [3], or by playing with the radial field one can reduce its divergence [4]. I'll then show how by controlling the quiver motion of relativistic electrons intense and bright X-rays beam are produced in a compact and elegant way [5-7]. Finally I'll show some examples of applications.

References

- [1] V. Malka, *Phys. of Plasmas* **19**, 055501 (2012)
- [2] E. Guillaume *et al.*, *Phys. Rev. Lett.* **115**, 155002 (2015).
- [3] C. Thauray *Scientific Report*, 10.1038, srep16310, Nov. 9 (2015)
- [4] C. Thauray *et al.*, *Nature Comm.* **6**, 6860 (2015)
- [5] K. Ta Phuoc *et al.*, *Nature Photonics* **6**, 308-311 (2012)
- [6] S. Corde *et al.*, *Review of Modern Phys.* **85** (2013)
- [7] I. Andriyash *et al.*, *Nature Comm.* **5**, 4736 (2014)